

Evaluation of greenhouse gas emissions from fertilized grassland



L. Balezientiene, E. Klimas, R. Bleizgys

Lithuanian University of Agriculture (LUA), Studentų 11, LT-53361, Akademija, Kaunas distr., Lithuania lgita.balezientiene@lzuu.lt



Agriculture contributes 10–12 % of the total global anthropogenic GHG emissions or about 6.8 Gt of CO₂ equivalent (eq) per year (UN FCCC, 2008). During 1990 - 2005, emissions from agriculture increased by about 17 % and are projected to increase further due to expected increases in food demand and diet changes as the global population continues to grow. Agricultural land occupied 5023 Mha in

Grassland (1.2 mln ha, or 50 % of agricultural lands in Lithuania) and fertilized areas occupy about 20% of the Earth's land surface generating considerable GHG emissions and increasing global climate change.

Objectives of this investigation were to determine impact of mono and complex fertilizers, their rates and combinations on

Methods. Field experiments were carried out on fertilized natural sward and cultural pasture at Training Farm of Lithuanian University of Agriculture (54.52 N, 23.50 E) Kaunas district, Calc(ar)-Endohypogleyic Luvisol, clay loam topsoil over silt loam. N and NPK application scheme in semi natural sward see Fig. 1. Cultural pasture was fertilized with N₁₈₀P₁₂₀K₁₅₀. P and K applied before vegetation in early spring, N fertilizer applied in 2 times: end of April and after 1st cut (beginning of July). GHG emissions were monitored by the chamber method ($r=0.43$ m, 0.05 m³), with 6 replicates. The measurements were done every 1-3 weeks between July and September in absence of frosts stress. The gas samples were analyzed in the laboratory by infrared gas analyzer (MGA3000). Daily net ecosystem CO₂, CH₄ and N₂O exchange ($\text{mg h}^{-1} \text{m}^{-2}$) were calculated by integrating the 60-minute fluxes determined by the meteorological measurements over each day.

Results




Figure 1. N₂O, CO₂ and CH₄ exchange between crops and atmosphere in natural sward (Control N₁₈₀P₁₂₀K₁₅₀) and cultural pasture (N₁₈₀P₁₂₀K₁₅₀) (mean±SE)




Figure 2. Correlation ($p < 0.05$) between N fertilizers rates and GHG emission

Conclusions. CO₂, N₂O and CH₄ emissions were nearly the same in natural and cultural grasslands and statistically significant correspond with fertilizing and abiotic environment factors. Employing natural sward would not to exceed N₆₀P₄₀K₅₀ fertilizing rate due to extensiveness of foraging species.

Acknowledgements. Research was carried in cooperation with ES COST ES0804 activity. This research was supported by the Agency for International Science and Technology Development Programmes in Lithuania during 2009.