

Grazing intensity and precipitation affects herbage accumulation, herbage quality, and animal performance in semiarid grassland

P. Schönbach, H. Wan, M. Gierus, K. Müller, L. Lin, A. Susenbeth, and F. Taube

Background

- Increasing grazing pressure on native grassland in Inner Mongolia, P.R. China
- Overgrazing and degradation (more than 80% of the typical steppe ecosystem in Inner Mongolia is degraded).
- Grazing not only alters the ecological functionality of the grassland but also its value for livestock farming.
- **Aim of the present study:** Analysis of grazing-induced changes in productivity and herbage quality as well as its effect on animal performance.



Fig. 1: Area of research, Inner Mongolia, P.R. China

Materials and methods

- Controlled grazing experiment (initiated in 2005)
- Experimental factors: Grazing intensity, Year (i.e. different rates of precipitation)
- Parameters:
 - a) Productivity: end-of-season herbage mass, herbage accumulation, live weight gain (LWG)
 - b) Herbage quality: e.g. crude protein, neutral detergent fibre, acid detergent fibre, acid detergent lignin, in-vitro digestibility, metabolizable energy
- Near-Infrared-Spectroscopy (NIRS) technique
- Statistics: ANOVA by using the Mixed Model, simple regression analysis in SAS version 9.1.

RESULTS (2005-2008)




Fig. 2: Relationship between precipitation and herbage accumulation (HA), crude protein (CP), in-vitro digestibility (CDOM), and neutral detergent fibre (NDF).




Fig. 3: Relationship between stocking rate and crude protein (CP), in-vitro digestibility (CDOM), neutral detergent fibre (NDF), and lignin (ADL).




Fig. 4: Relationship between stocking rate and herbage accumulation (HA).
 (HA [g DM/m²] = -9.6 × SR + 209.9, r² = 0.3*** RMSE = 43.9)




Fig. 5: Relationship between stocking rate and live-weight-gain (LWG).
 LWG per sheep (g LW/d) = -0.6369x² + 3.7919x + 84.5045, r² = 0.32** RMSE = 14.5, peak: 3 sheep/ha
 LWG per ha (g LW/d) = -4.2805x² + 109.5529x - 25.8257, r² = 0.8*** RMSE = 86.7, peak: 13 sheep/ha

Conclusions

- Precipitation appeared to be the crucial factor determining accumulation and quality of herbage.
- Grazing intensity played the key role in determining animal performance.
- Increasing grazing intensity decreased herbage accumulation but increased herbage quality.
- Animal performance in this rangeland was primarily influenced by herbage quantity. There were no benefits of grazing-induced increases in herbage quality on live-weight-gain.

