

White clover effect on yield and quality of a *Lolium perenne* sward under cutting conditions

De Vliegheer A.

ILVO Plant Sciences Unit, 9820 Merelbeke, Belgium

Introduction

Farmers in Flanders see great potential in introducing white clover (WC) into their grassland, because N-fertilisation is expensive and restricted by the Flemish Government in accordance with the EU Nitrate Directive. Is there any effect of WC in the perennial ryegrass sward when 200-330 N/ha is applied? Are there consequences for the leachable N residue in the soil?

Materials and methods

- Experiment: 2007-2011
- Sandy loam soil in Merelbeke (Belgium)
- Split plot design - harvest area per plot: 8.4 m² - 5 cuts a year.
- Sward types:
 - perennial ryegrass (PRG)
 - perennial ryegrass + white clover (PRG/WC)
- Fertilisation treatments
 - (1) 170 N_{slurry} + 100 N_{mineral} (2) 170 N_{slurry} + 180 N_{mineral}
 - (3) 250 N_{slurry} + 100 N_{mineral} (4) 250 N_{slurry} + 180 N_{mineral}



Measurements:

- Dry matter yield, clover content, forage quality
- Calculation of energy value and protein quality
- Nitrate nitrogen content of the 0-90 cm top soil in autumn

Results

	2007		2008	
	PRG	PRG/WC	PRG	PRG/WC
Yield (ha ⁻¹ y ⁻¹)				
Dry matter (kg)	13516 ^a	13099 ^a	13284 ^a	14305 ^b
Crude protein (kg)	2090 ^a	2130 ^a	2297 ^a	2684 ^b
Clover content in DM (g kg ⁻¹)	-	130	-	210
Forage quality				
Energy VEM in DM (kg ⁻¹)	848 ^a	837 ^a	862 ^a	859 ^a
Crude protein in DM (g kg ⁻¹)	155 ^a	163 ^b	183 ^a	200 ^b
DVE in DM (g kg ⁻¹)	77 ^a	78 ^a	84 ^a	89 ^b
OEB in DM (g kg ⁻¹)	3 ^a	12 ^a	26 ^a	40 ^a
Residual nitrate in soil				
NO ₃ -N (kg ha ⁻¹)	15 ^a	24 ^b	6 ^a	11 ^b

VEM: fodder unit milk; DVE: true protein digested in the small intestine; OEB: rumen degraded protein balance
Treatments with the same letter in the same row and in the same year are not significantly different (P<0.05)

Conclusions

In a cutting regime with N-fertilisation between 200 and 330 N_{available} ha⁻¹ white clover is still an interesting component in a perennial ryegrass sward because:

- it can persist in the sward
- it can stimulate DM and CP yield and increase CP content of the forage
- it increases the nitrate residue in the soil significantly without any risk for nitrate leaching (values far below the threshold of 90 kg NO₃-N ha⁻¹ in 0-90 cm top soil)



www.ilvo.vlaanderen.be

