

# A simple model for the estimation of protein content of first-cut meadow forage

Peratoner G.<sup>1</sup>, Bodner A.<sup>1</sup>, Stimpfl E.<sup>1</sup>, Werth E.<sup>1</sup>, Schaumberger A.<sup>2</sup>, Kasal A.<sup>1</sup>

<sup>1</sup> Research Centre for Agriculture and Forestry Laimburg

<sup>2</sup> AREC Raumberg-Gumpenstein


## Introduction

- Forage quality of meadows declines over time with increasing developmental stage of plants; harvest timing is pivotal to the achievement of satisfactory forage quality
- 5-year data were collected in South Tyrol with the aim of developing an affordable and reliable tool for estimating forage quality as a function of harvest timing and other factors related to vegetation, geomorphology, soil properties and agronomic management
- Crude protein data (CP) have been chosen as a dependent variable for model development
- Focus of the present paper: replacing year effects, not useful for predictive purposes, with other factors related to the climatic features of the sites and to the weather conditions during vegetation growth

## Material and methods

- 35 sampling sites, 5 years
- Weekly sequential sampling starting at pasture stage (15 cm growing height) – see poster 3.1.36 for details
- Statistical analysis: binary logistic regression
  - dependent dichotomous variable: CP < 150 g kg<sup>-1</sup> or CP ≤ 150 g kg<sup>-1</sup>
  - classification cut off 0.5

Altitude range: 667 - 1593 m a.s.l.  
Cutting regime: 2 to 5 cuts y<sup>-1</sup>



### FACTORS GROUP A

**Harvest date**  
• Days since pasture stage

**Soil properties**  
• texture  
• organic matter  
• pH  
• P  
• K

**Agronomic management**  
• cut frequency  
• N-input  
• dung type  
• irrigation

**Botanical composition**  
• Functional groups dominance  
• Vegetation type




**Weather conditions**  
• Agricultural district  
• Year

**Geomorphologic traits**  
• Altitude  
• Slope  
• Exposition

### FACTORS GROUP B

Sum and mean of daily temperature departures from long-term average between 1 week before pasture stage and harvest date (reference weather stations of the respective agricultural district)

Solar radiation sum between 1 week before pasture stage and harvest date



## Results

- Analysis 1 (factors group A): **10 significant factors, mean correct prediction 87.5%**
- Analysis 2 (replacement of year and time since pasture stage with factors group B): **15 significant factors, mean correct prediction 86.8%**

## Conclusions

- Factors including cumulative and/or average values of the potential solar radiation, and of the temperature departure from the long-term average for the respective agricultural district, successfully replaced the year effect in the logistic regression and resulted in a satisfactory prediction accuracy

## Contact

Dr. Giovanni Peratoner – Section Mountain Agriculture  
Land- und Forstwirtschaftliches Versuchszentrum Laimburg | Centro di Sperimentazione Agraria e Forestale Laimburg  
Research Centre for Agriculture and Forestry Laimburg | Laimburg 6 – Pfatten (Vadena) | 39040 Auer (Ora) | Südtirol (Alto Adige) | Italy