

A scheduling model for forage harvesting

Claus Grøn Sørensen
Dionysis Bochtis
Ole Green
Frank Oudshoorn



Background



Grass silage quality has major influence on milk yield

Grass silage quality can save import of concentrates

It has influence on the GHG budget.

In Denmark 1 /3 of grass silage is too wet and 1 /3 is too dry.

Many planning models on harvest and route planning have been launched

Objective

Can we make a scheduling model for the optimal time for initiating various operations related to the harvesting and treatment of grass forage.

The scheduling model is based on moisture predictions

The model anticipates three levels of decision:

1. Harvest
2. Uncertainty
3. No harvest

minimum time lags → (time window) ← maximum time lags



biomass operations

Inputs for the model

Should be possible for the farmer to fill-in

Expected yield (would prefer biomass map)

Potential field working hours for cut

Potential field working hours for picking up


Potential field working hours for spreading and raking

Field coverage

Moisture prediction model of Atzema, (can use online weather forecast)

The DSS


1st stage decision making (1sDM)



The DSS






FACULTY OF AGRICULTURAL SCIENCES
AARHUS UNIVERSITY



The Moisture prediction model (MPM)



FACULTY OF AGRICULTURAL SCIENCES
AARHUS UNIVERSITY




Average absolute error
= 2 percent

The DSS

1st stage decision making (1sDM)



FACULTY OF AGRICULTURAL SCIENCES
AARHUS UNIVERSITY



The distribution of the predicted levels of moisture content by the 1sDM


The percentage of successful predictions

The DSS

2nd stage decision making (2sDM)



FACULTY OF AGRICULTURAL SCIENCES
AARHUS UNIVERSITY




Different covering percentage results in different drying rates for the grass

The DSS

2nd stage decision making (2sDM)



FACULTY OF AGRICULTURAL SCIENCES
AARHUS UNIVERSITY





The DSS

2nd stage decision making (2sDM)



FACULTY OF AGRICULTURAL SCIENCES
AARHUS UNIVERSITY




The distribution of the predicted levels of moisture content by the 2sDM

The percentage of successful predictions

DSS evaluation



FACULTY OF AGRICULTURAL SCIENCES
AARHUS UNIVERSITY



Unwanted outcomes from harvesting decision making based on unsuccessful level predictions

DSS evaluation

Percentages of the undesirable outcomes from unsuccessful level predictions

The farmer proceeds to the harvesting operation based on the decision 2 suggestion

& The farmer does not proceed to the harvesting operation based on the decision 2 suggestion

		reduced quality (%)	poor quality (%)	lost opportunity (%)		Total (%)
				acceptable quality	high quality	
1sDM	D2+#	1.99	3.12	0.28	0	5.39
	D2-&	1.99	0.28	0.28	12.22	14.77
2sDM	D2+	4.26	6.54	0	0.57	11.37
	D2-	4.26	0.57	0	1.14	5.97


Conclusion

- The percentage of successful suggestions ranged from 83.33% to 94.61%.
- The developed model may be seen as a prototype for a decision support system, providing recommendations on the optimal execution of grass harvesting operations.
- The development of such a system in a contractor-level (multiple-fields) is necessary and could provide multiple benefits.

scheduling tool for biomass field logistics



FACULTY OF AGRICULTURAL SCIENCES
AARHUS UNIVERSITY



Thank you for your attention !

